

## Answer to Some Selected Problems

### UNIT 8

8.25 15 g

### UNIT 12

12.32 Mass of carbon dioxide formed = 0.505 g

Mass of water formed = 0.0864 g

12.33 % of nitrogen = 56

12.34 % of chlorine = 37.57

12.35 % of sulphur = 19.66

### UNIT 13

13.1 Due to the side reaction in termination step by the combination of two  $\dot{\text{C}}\text{H}_3$  free radicals.

- 13.2 (a) 2-Methyl-but-2-ene (b) Pent-1-ene-3-yne  
 (c) Buta-1, 3-diene (d) 4-Phenylbut-1-ene  
 (e) 2-Methylphenol (f) 5-(2-Methylpropyl)-decane  
 (g) 4-Ethyldeca -1,5,8- triene


- 13.3 (a) (i)  $\text{CH}_2 = \text{CH} - \text{CH}_2 - \text{CH}_3$  But-1-ene  
 (ii)  $\text{CH}_3 - \text{CH}_2 = \text{CH} - \text{CH}_3$  But-2-ene  
 (iii)  $\text{CH}_2 = \underset{\text{CH}_3}{\text{C}} - \text{CH}_3$  2-Methylpropene
- (b) (i)  $\text{HC} \equiv \text{C} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$  Pent-1-yne  
 (ii)  $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH}_3$  Pent-2-yne  
 (iii)  $\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \text{C} \equiv \text{CH}$  3-Methylbut-1-yne

- 13.4 (i) Ethanal and propanal (ii) Butan-2-one and pentan-2-one  
 (iii) Methanal and pentan-3-one (iv) Propanal and benzaldehyde


13.5 3-Ethylpent-2-ene

13.6 But-2-ene

13.7 4-Ethylhex-3-ene



- 13.8 (a)  $C_4H_{10}(g) + 13/2 O_2(g) \xrightarrow{\Delta} 4CO_2(g) + 5H_2O(g)$   
 (b)  $C_5H_{10}(g) + 15/2 O_2(g) \xrightarrow{\Delta} 5CO_2(g) + 5H_2O(g)$   
 (c)  $C_6H_{10}(g) + 17/2 O_2(g) \xrightarrow{\Delta} 6CO_2(g) + 5H_2O(g)$   
 (d)  $C_7H_8(g) + 9O_2(g) \xrightarrow{\Delta} 7CO_2(g) + 4H_2O(g)$




The *cis* form will have higher boiling point due to more polar nature leading to stronger intermolecular dipole-dipole interaction, thus requiring more heat energy to separate them.

13.10 Due to resonance


13.11 Planar, conjugated ring system with delocalisation of  $(4n+2)$   $\pi$  electrons, where,  $n$  is an integer

13.12 Lack of delocalisation of  $(4n+2)$   $\pi$  electrons in the cyclic system.




13.13 (i)




(ii)



(iii)



13.14




15 H attached to 1° carbons

4 H attached to 2° carbons

1 H attached to 3° carbons

13.15 More the branching in alkane, lower will be the boiling point.





13.16 Refer to addition reaction of HBr to unsymmetrical alkenes in the text.



All the three products cannot be obtained by any one of the Kekulé's structures. This shows that benzene is a resonance hybrid of the two resonating structures.

 13.18  $\text{H} - \text{C} \equiv \text{C} - \text{H} > \text{C}_6\text{H}_6 > \text{C}_6\text{H}_{14}$ . Due to maximum s orbital character in ethyne (50 per cent) as compared to 33 per cent in benzene and 25 per cent in n-hexane.

13.19 Due to the presence of 6 π electrons, benzene behaves as a rich source of electrons thus being easily attacked by reagents deficient in electrons.



13.22 (a) Chlorobenzene > *p*-nitrochlorobenzene > 2,4 - dinitrochlorobenzene

(b) Toluene > *p*-CH<sub>3</sub>-C<sub>6</sub>H<sub>4</sub>-NO<sub>2</sub> > *p*-O<sub>2</sub>N-C<sub>6</sub>H<sub>4</sub>-NO<sub>2</sub>

13.23 Toluene undergoes nitration most easily due to electron releasing nature of the methyl group.

13.24 FeCl<sub>3</sub>

13.25 Due to the formation of side products. For example, by starting with 1-bromopropane and 1-bromobutane, hexane and octane are the side products besides heptane.

----- Click on the -----  
**Download Link**  
 To view the complete book

## INDEX

### A

Acid rain	409, 410		
Acidic dehydration of alcohols	388		
Activating groups	403		
Acyclic compounds	339		
Alicyclic compounds	339		
Alkali metals	300		
– atomic radii	300		
– chemical properties	301		
– halides	303		
– hydration enthalpy	300		
– hydroxides	303		
– ionic radii	300		
– ionisation enthalpy	300		
– oxides	303		
– physical properties	300		
– reactivity towards air	301		
– reactivity towards dihydrogen	302		
– reactivity towards halogens	302		
– reactivity towards water	302		
– reducing nature	302		
– salts of oxoacids	303		
– solution in liquid ammonia	302		
– uses	302		
Alkaline earth metals	306		
– atomic radii	306		
– carbonates	309		
– chemical properties	308		
– electronic configuration	306		
– halides	309		
– hydration enthalpies	307		
– hydroxides	309		
– ionic radii	306		
– ionisation enthalpies	307		
– nitrates	309		
– oxides	309		
– physical properties	307		
– reactivity towards air	308		
– reactivity towards halogens	308		
– reactivity towards water	308		
– reducing nature	308		
– salts of oxoacids	309		
– solution in liquid ammonia	308		
– uses	308		
– sulphates	309		
Alkanes	374		
– aromatisation	382		
– chain isomerism	386		
– combustion	381		
– controlled oxidation	382		
– geometrical isomerism	386		
– halogenation	380		
		– halogenation mechanism	381
		– Isomerisation	382
		– ozonolysis	391
		– pyrolysis	382
		– reaction with steam	382
		Alkenes	384
		– addition of dihydrogen	388
		– addition of hydrogen halides	389
		– addition of hydrogen halides, mechanism	389
		– addition of sulphuric acid	390
		– addition of water	391
		– chemical properties	388
		– geometrical isomers	386
		– oxidation	391
		– physical properties	389
		– position isomerism	386
		– preparation	387
		– structural isomerism	385
		Alkynes	392
		– acidic characters	394
		– addition of dihydrogen	395
		– addition of halogens	395
		– addition of hydrogen halides	395
		– addition of water	395
		– addition reaction	394
		– cyclic polymerisation	396
		– linear polymerisation	395
		– polymerisation	395
		– preparation	393
		Allotropes of carbon	325
		Aluminium	317, 318, 322
		Aluminium, uses	322
		Angle of torsion	384
		Anti Markovnikov rule	390
		Arenes	396
		Arenium ion, formation	401
		Arenium ion, stabilisation	401
		Aromatic compounds	339
		Aromaticity	399
		Atmospheric pollution	407
		<b>B</b>	
		Baking soda	306
		Balancing of redox reaction	274
		Benzene	396
		– Friedel-Crafts alkylation	400
		– chemical properties	400
		– combustion	402
		– electrophilic substitution	400
		– Friedel-Crafts acylation	400
		– mechanism of electrophilic substitution	401
		– nitration	400